

## 2.6 Domain and Range

## 2.6 Domain and Range

### Learning Targets:


1. SWBAT determine the domain and range of a function from its graph.
2. SWBAT determine the domain and range of a function from its equation.




**Domain** of a function is the set of **x values** for which a value of  $y$  can be determined.

**Range** of a function consists of all the **y values** of a function.



# Determining Domain and Range from the Graph of a Function



$$D: (-\infty, \infty)$$
$$R: (-\infty, \infty)$$



$$D: (-\infty, \infty)$$
$$R: [-2, \infty)$$



$$D: (-\infty, \infty)$$

$$R: (-\infty, 0) \cup [2, \infty)$$

$$R: [-2, 0]$$

# Determining the Domain By Examining The Function

To find the domain of a function without seeing the graph we must be careful to exclude any value of  $x$  that will result in:

Division by 0.

Taking the even root of a negative number.

Finding the logarithm of a non-positive number.

A trig function being undefined

Ex.1 Determine the domain of the following functions:

a)  $f(x) = x^6 - 15x^3$

polynomial

$D: (-\infty, \infty)$



Rational

$$\text{b) } f(x) = \frac{10}{x^3 - 9x}$$

$$= \frac{10}{x(x^2 - 9)} = \frac{10}{x(x-3)(x+3)}$$

$$x \neq 0, 3, -3$$


$$(-\infty, -3) \cup (-3, 0) \cup (0, 3) \cup (3, \infty)$$



$$c) f(x) = \sqrt{x^2 + x - 30}$$

$$x^2 + x - 30 \geq 0$$

$$(x+6)(x-5) \geq 0$$



$$D: (-\infty, -6] \cup [5, \infty)$$

$$\sqrt{x^2 + x - 30}$$
$$(-\infty, -6) \cup (5, \infty)$$


$$d) h(x) = \log_5(2x - 7)$$



$$2x - 7 > 0$$

$$2x > 7$$

$$x > 7/2$$



$$e) g(x) = \sqrt[3]{\frac{2}{x} - 4}$$

D:

$$(-\infty, 0) \cup (0, \infty)$$

OR

$$x \neq 0$$

## Assignment Page 99

#'s 1-12 (Domain and Range Off of Graph)

#'s 13-21,27,31,33,35,37,38,42 (just find domain algebraically)

## Determining the Range By Examining The Function

This is harder to do. Later in our course we will develop strategies that will assist us. For now, we need to use a good deal of common sense and past knowledge.

Ex.2 Determine the range of the following functions:

a)  $f(x) = x^3 - 9x$

D:  $(-\infty, \infty)$

$$\frac{\quad}{S} \quad \frac{\quad}{E}$$

R:  $(-\infty, \infty)$

\* Any odd degree polynomial has all the reals as domain/range.

$$b) f(x) = (x^2 + 5)^2$$

$$D: (-\infty, \infty)$$

$$R: [25, \infty)$$

x	y
0	25
1	36
-1	36
2	81
-2	81

$$c) f(x) = \sqrt{4+x}$$

$$= \sqrt{x+4}$$


$$D: [-4, \infty)$$

$$R: [0, \infty)$$





$$d) f(x) = \sqrt{100 - x^2}$$



$$D: [-10, 10]$$

$$R: [0, 10]$$

$$x^2 + y^2 = r^2$$


$$y^2 = r^2 - x^2$$

$$y = \pm \sqrt{r^2 - x^2}$$

$$f(x) = \sqrt{100 - x^2}$$

$$D: 100 - x^2 \geq 0$$

$$(10 - x)(10 + x) \geq 0$$



x	y
-10	0
0	10
10	0

$$e) f(x) = \sqrt[3]{x-1}$$

$$D: (-\infty, \infty)$$


$$R: (-\infty, \infty)$$

$$f) f(x) = \frac{3x^0}{x^1 - 2}$$

$$x \neq 2$$

$$D: (-\infty, 2) \cup (2, \infty)$$

$$R: (-\infty, 0) \cup (0, \infty)$$




$$g) f(x) = \frac{4}{x^2 + 2}$$

$$D: (-\infty, \infty)$$

$$R: (0, 2]$$

(0, 2)



x	y
0	2
1	4/3
-1	4/3
-2	2/3
2	2/3

↓ ↓

$$f(-5) = (-5)^2 + 10(-5) \\ = -25$$

$$h) f(x) = x^2 + 10x$$

$$y = a(x-h)^2 + q$$


$$D: (-\infty, \infty)$$

$$R: [-25, \infty)$$

$$x^2 + 10x = 0$$

$$x(x+10) = 0$$

$$x = 0 \quad x = -10$$



$$f(x) = x^2 + 10x$$

$$= \underbrace{x^2 + 10x + 25}_{(x+5)^2} - \underline{25}$$

$$f(x) = (x+5)^2 - 25$$

### **TO FIND THE RANGE, CONSIDER THESE QUESTIONS**

- Is the function a translation of a function whose graph I already know?
- Is it a polynomial function? If so examine the degree and the leading coefficient.
- What happens to  $y$  as  $x$  becomes very large positive (negative)?
- What happens to  $y$  near any vertical asymptotes? Use a sign analysis.
- Is there a horizontal asymptote line?
- What is the  $y$ -intercept?
- If you are dealing with a root function, is there are largest (smallest) value for the expression beneath the radical sign?

Page 98



Assignment Page 99

#'s 13-21,27,31,33,35,37,38,42 (just find range algebraically)