

5.3 Definite Integrals and Antiderivatives

5.3 Definite Integrals and Anti-Derivatives

Learning Targets:

1. SWBAT use the properties of definite integrals to evaluate definite integrals.
2. SWBAT to use definite integral rules to manually evaluate definite integrals.



Properties of Definite Integrals

1. *Order of Integration:* $\int_b^a f(x) dx = -\int_a^b f(x) dx$ A definition

2. *Zero:* $\int_a^a f(x) dx = 0$ Also a definition

3. *Constant Multiple:* $\int_a^b kf(x) dx = k \int_a^b f(x) dx$ Any number k

$$\int_a^b -f(x) dx = -\int_a^b f(x) dx \quad k = -1$$

4. *Sum and Difference:* $\int_a^b (f(x) \pm g(x)) dx = \int_a^b f(x) dx \pm \int_a^b g(x) dx$

5. *Additivity:* $\int_a^b f(x) dx + \int_b^c f(x) dx = \int_a^c f(x) dx$

21. Given $\int_0^5 f(x) dx = 10$ and $\int_5^7 f(x) dx = 3$, find

(a) $\int_0^7 f(x) dx.$ (b) $\int_5^0 f(x) dx.$

(c) $\int_5^5 f(x) dx.$ (d) $\int_0^5 3f(x) dx.$

22. Given $\int_0^3 f(x) dx = 4$ and $\int_3^0 f(x) dx = -1$, find

(a) $\int_0^6 f(x) dx.$ (b) $\int_6^3 f(x) dx.$

(c) $\int_3^3 f(x) dx.$ (d) $\int_3^6 -5f(x) dx.$

23. Given $\int_2^6 f(x) dx = 10$ and $\int_2^6 g(x) dx = -2$, find

(a) $\int_2^6 [f(x) + g(x)] dx.$ (b) $\int_2^6 [g(x) - f(x)] dx.$

(c) $\int_2^6 2g(x) dx.$ (d) $\int_2^6 3f(x) dx.$

24. Given $\int_{-1}^1 f(x) dx = 0$ and $\int_0^1 f(x) dx = 5$, find

(a) $\int_{-1}^0 f(x) dx.$ (b) $\int_0^1 f(x) dx - \int_{-1}^0 f(x) dx.$

(c) $\int_{-1}^1 3f(x) dx.$ (d) $\int_0^1 3f(x) dx.$

Basic Integration Rules

$$\int c dx = cx$$

$$\int x^n dx = \frac{x^{n+1}}{n+1}, n \neq -1$$

$$\int \cos x dx = \sin x$$

$$\int \sin x dx = -\cos x$$

$$\int \frac{1}{x} dx = \ln|x|$$

$$\int e^x dx = e^x$$

$$\int e^{kx} dx = \frac{e^{kx}}{k}$$

$$\int a^{kx} dx = \frac{a^{kx}}{k \ln a}$$

Ex.1 Evaluate the following definite integrals:

$$\text{a) } \int_3^5 2x dx$$

$$\text{b) } \int_1^4 (2x + 3) dx$$

$$\text{c) } \int_{\pi/3}^{\pi/6} \cos x dx$$

$$\text{d) } \int_{\pi/30}^{\pi/10} \sin 5x dx$$

$$\text{e) } \int_{\ln 4}^{\ln 6} e^x dx$$

$$f) \int_5^{10} \frac{1}{s} ds$$

$$\text{g) } \int_1^3 \frac{w^2 + 1}{w} dw$$

#8 from handout

2003 MC Question

2. $\int_0^1 e^{-4x} dx =$

(A) $\frac{-e^{-4}}{4}$

(B) $-4e^{-4}$

(C) $e^{-4} - 1$

(D) $\frac{1}{4} - \frac{e^{-4}}{4}$

(E) $4 - 4e^{-4}$

Example

Let f be a continuous function such that $\int_8^0 f(x) dx = -13$ and $\int_0^2 f(x) dx = 8$. What is the value of $\int_2^8 (f(x) + 4) dx$?

<http://archives.math.utk.edu/visual.calculus/4/ftc.10/index.html>

**AP[®] CALCULUS AB
2005 SCORING GUIDELINES**

Question 2

The tide removes sand from Sandy Point Beach at a rate modeled by the function R , given by

$$R(t) = 2 + 5 \sin\left(\frac{4\pi t}{25}\right).$$

A pumping station adds sand to the beach at a rate modeled by the function S , given by

$$S(t) = \frac{15t}{1 + 3t}.$$

Both $R(t)$ and $S(t)$ have units of cubic yards per hour and t is measured in hours for $0 \leq t \leq 6$. At time $t = 0$, the beach contains 2500 cubic yards of sand.

- (a) How much sand will the tide remove from the beach during this 6-hour period? Indicate units of measure.
- (b) Write an expression for $Y(t)$, the total number of cubic yards of sand on the beach at time t .
- (c) Find the rate at which the total amount of sand on the beach is changing at time $t = 4$.
- (d) For $0 \leq t \leq 6$, at what time t is the amount of sand on the beach a minimum? What is the minimum value?
Justify your answers.

AP[®] CALCULUS AB
2005 SCORING GUIDELINES (Form B)

Question 2

A water tank at Camp Newton holds 1200 gallons of water at time $t = 0$. During the time interval $0 \leq t \leq 18$ hours, water is pumped into the tank at the rate

$$W(t) = 95\sqrt{t} \sin^2\left(\frac{t}{6}\right) \text{ gallons per hour.}$$

During the same time interval, water is removed from the tank at the rate

$$R(t) = 275 \sin^2\left(\frac{t}{3}\right) \text{ gallons per hour.}$$

- (a) Is the amount of water in the tank increasing at time $t = 15$? Why or why not?
- (b) To the nearest whole number, how many gallons of water are in the tank at time $t = 18$?
- (c) At what time t , for $0 \leq t \leq 18$, is the amount of water in the tank at an absolute minimum? Show the work that leads to your conclusion.
- (d) For $t > 18$, no water is pumped into the tank, but water continues to be removed at the rate $R(t)$ until the tank becomes empty. Let k be the time at which the tank becomes empty. Write, but do not solve, an equation involving an integral expression that can be used to find the value of k .

AP[®] CALCULUS AB
2004 SCORING GUIDELINES (Form B)

Question 2

For $0 \leq t \leq 31$, the rate of change of the number of mosquitoes on Tropical Island at time t days is modeled by $R(t) = 5\sqrt{t} \cos\left(\frac{t}{5}\right)$ mosquitoes per day. There are 1000 mosquitoes on Tropical Island at time $t = 0$.

- (a) Show that the number of mosquitoes is increasing at time $t = 6$.
- (b) At time $t = 6$, is the number of mosquitoes increasing at an increasing rate, or is the number of mosquitoes increasing at a decreasing rate? Give a reason for your answer.
- (c) According to the model, how many mosquitoes will be on the island at time $t = 31$? Round your answer to the nearest whole number.
- (d) To the nearest whole number, what is the maximum number of mosquitoes for $0 \leq t \leq 31$? Show the analysis that leads to your conclusion.

Assignment

AP Calc Text Page 274

#'s 1-12, 15

Calc 30 Text Page 370

#'s 1-20, 39

Average Value of a Function

Average Value Video

Average Value of a Function on an Interval

If f is a continuous function on $[a,b]$, then the **Average Value** of f on $[a,b]$ is:


$$= \frac{1}{b-a} \int_a^b f(x) dx$$

Example 1

Find the average value of $y = \sin x$ between $x = 0$ and $x = \pi$.

Example 2


The graph of f is shown, find the average value of f on $[0,4]$.



Mean Value Theorem for Integrals

If f is a continuous function on $[a,b]$, then at some point c in $[a,b]$:

$$f(c)(b-a) = \int_a^b f(x) dx$$



Mean Value Theorem

Example 3

Given $f(x) = \sqrt{x-1}$, verify the hypothesis of the Mean Value Theorem for Integrals for f on $[1, 10]$ and find the value of c indicated in the theorem.

Example: Given that the velocity function for a given object by $v(t) = 3t^2 - 18t + 24$, where t is given in seconds and $v(t)$ is in m/s. Find the average velocity between $t=1$ s and $t=3$ s.






2013 AP[®] CALCULUS AB FREE-RESPONSE QUESTIONS**CALCULUS AB****SECTION II, Part B****Time—60 minutes****Number of problems—4****No calculator is allowed for these problems.**

t (minutes)	0	1	2	3	4	5	6
$C(t)$ (ounces)	0	5.3	8.8	11.2	12.8	13.8	14.5

3. Hot water is dripping through a coffeemaker, filling a large cup with coffee. The amount of coffee in the cup at time t , $0 \leq t \leq 6$, is given by a differentiable function C , where t is measured in minutes. Selected values of $C(t)$, measured in ounces, are given in the table above.
- (a) Use the data in the table to approximate $C'(3.5)$. Show the computations that lead to your answer, and indicate units of measure.
- (b) Is there a time t , $2 \leq t \leq 4$, at which $C'(t) = 2$? Justify your answer.
- (c) Use a midpoint sum with three subintervals of equal length indicated by the data in the table to approximate the value of $\frac{1}{6} \int_0^6 C(t) dt$. Using correct units, explain the meaning of $\frac{1}{6} \int_0^6 C(t) dt$ in the context of the problem.
- (d) The amount of coffee in the cup, in ounces, is modeled by $B(t) = 16 - 16e^{-0.4t}$. Using this model, find the rate at which the amount of coffee in the cup is changing when $t = 5$.

88. On the closed interval $[2, 4]$, which of the following could be the graph of a function f with the property that

$$\frac{1}{4-2} \int_2^4 f(t) dt = 1?$$



Things Not Said By Calculus Students

Assignment

Page 275

#'s 21-24 as a group

#'s 25-30